

COHOMOLOGY RING OF DIFFERENTIAL OPERATOR RINGS

GRACIELA CARBONI, JORGE A. GUCCIONE, AND JUAN J. GUCCIONE

ABSTRACT. We compute the multiplicative structure in the Hochschild cohomology ring of a differential operators ring and the cap product of Hochschild cohomology on the Hochschild homology.

INTRODUCTION

Let k be a field and A an associative k -algebra with 1. An extension E/A of A is a *differential operator ring* on A if there exists a Lie k -algebra $\mathfrak{g}$ and a k -vector space embedding $x \mapsto \bar{x}$, of $\mathfrak{g}$ into E , such that for all $x, y \in \mathfrak{g}$, $a \in A$:

- (1) $\bar{a}a - a\bar{a} = a^x$, where $a \mapsto a^x$ is a derivation,
- (2) $\overline{xy} - \overline{yx} = \overline{[x, y]_{\mathfrak{g}}} + f(x, y)$, where $[-, -]_{\mathfrak{g}}$ is the bracket of $\mathfrak{g}$ and $f: \mathfrak{g} \times \mathfrak{g} \rightarrow A$ is a k -bilinear map,
- (3) for a given basis $(x_i)_{i \in I}$ of $\mathfrak{g}$, the algebra E is a free left A -module with the standard monomials in the x_i 's as a basis.

This general construction was introduced in [Ch] and [Mc-R]. Several particular cases of this type of extensions have been considered previously in the literature. For instance:

- when $\mathfrak{g}$ is one dimensional and f is trivial, E is the Ore extension $A[x, \delta]$, where $\delta(a) = a^x$,
- when $A = k$, one obtain the algebras studied by Sridharan in [S], which are the quasi-commutative algebras E , whose associated graded algebra is a symmetric algebra,
- in [Mc, §2] this type of extensions was studied under the hypothesis that A is commutative and $(x, a) \mapsto a^x$ is an action, and in [B-G-R, Theorem 4.2] the case in which the cocycle is trivial was considered.

In [B-C-M] and [D-T] the study of the crossed products $A \#_f H$ of a k -algebra A by a Hopf k -algebra H was begun, and in [M] was proved that the differential operator rings on A are the crossed products of A by enveloping algebras of Lie algebras.

In [G-G1] complexes, simpler than the canonical ones, giving the Hochschild homology and cohomology of a differential operator ring E with coefficients in an E -bimodule M , were obtained. In this paper we continue this investigation by studying the Hochschild cohomology ring of E and the cap product

$$H_p(E, M) \times HH^q(E) \rightarrow H_{p-q}(E, M) \quad (q \leq p),$$

2000 *Mathematics Subject Classification.* Primary 16E40; Secondary 16S32.

Key words and phrases. Differential operator rings; Hochschild (co)homology; cup product; cap product.

Supported UBACYT 095.

UBACYT 095 and PIP 112-200801-00900 (CONICET).

UBACYT 095 and PIP 112-200801-00900 (CONICET).

in terms of the above mentioned complexes. Moreover we generalize the results of [G-G1] by considering the (co)homology of E relative to a subalgebra K of A which is stable under the action of $\mathfrak{g}$ (which we also call the Hochschild (co)homology of the K -algebra E), and we seized the opportunity to fix some minor mistakes and to simplify some proofs in [G-G1].

The paper is organized in the following way: In Section 1 we obtain a projective resolution (X_*, d_*) of the E -bimodule E , relative to the family of all epimorphism of E -bimodules which split as (E, K) -bimodule maps. In Section 2 we determine and study comparison maps between (X_*, d_*) and the relative to K normalized Hochschild resolution $(E \otimes_K \overline{E}^{\otimes_K^*} \otimes_K E, b'_*)$ of E . In Sections 3 and 4 we apply the above results in order to obtain complexes $(\overline{X}_*^K(M), \overline{d}_*)$ and $(\overline{X}_K^*(M), \overline{d}^*)$, simpler than the canonical ones, giving Hochschild homology and cohomology of the K -algebra E with coefficients in an E -bimodule M , respectively. The main results are Theorems 3.4 and 4.4, in which we obtain morphisms

$$\overline{X}_K^*(E) \otimes \overline{X}_K^*(E) \rightarrow \overline{X}_K^*(E) \quad \text{and} \quad \overline{X}_*^K(M) \otimes \overline{X}_K^*(E) \rightarrow \overline{X}_*^K(M),$$

inducing the cup and cap product, respectively. Finally in Section 5, assuming that A is a symmetric algebra, we obtain further simplifications.

1. PRELIMINARIES

Let k be a field. In this paper all the algebras are over k . Let A be an algebra and H a Hopf algebra. We are going to use the Sweedler notation $\Delta(h) = \sum_{(h)} h^{(1)} \otimes_k h^{(2)}$ for the comultiplication Δ of H . A *weak action* of H on A is a k -bilinear map $(h, a) \mapsto a^h$, from $H \times A$ to A , such that

- (1) $(ab)^h = \sum_{(h)} a^{h^{(1)}} b^{h^{(2)}}$,
- (2) $1^h = \epsilon(h)1$,
- (3) $a^1 = a$,

for $h \in H$, $a, b \in A$. By an *action* of H on A we mean a weak action such that

$$(a^l)^h = a^{hl} \quad \text{for all } h, l \in H, a \in A.$$

Let A be an algebra and let H be a Hopf algebra acting weakly on A . Given a k -linear map $f: H \otimes_k H \rightarrow A$ we let $A \#_f H$ denote the algebra (in general non associative and without 1) whose underlying vector space is $A \otimes_k H$ and whose multiplication is given by

$$(a \otimes_k h)(b \otimes_k l) = \sum_{(h)(l)} ab^{h^{(1)}} f(h^{(2)}, l^{(1)}) \otimes_k h^{(3)} l^{(2)},$$

for all $a, b \in A$, $h, l \in H$. The element $a \otimes_k h$ of $A \#_f H$ will usually be written $a \# h$. The algebra $A \#_f H$ is called a *crossed product* if it is associative with $1 \# 1$ as identity element. In [B-C-M] it was proven that this happens if and only if the map f and the weak action of H on A satisfy the following conditions

- (1) (Normality of f) for all $h \in H$ we have $f(h, 1) = f(1, h) = \epsilon(h)1_A$,
- (2) (Cocycle condition) for all $h, l, m \in H$ we have

$$\sum_{(h)(l)(m)} f(l^{(1)}, m^{(1)})^{h^{(1)}} f(h^{(2)}, l^{(2)} m^{(2)}) = \sum_{(h)(l)} f(h^{(1)}, l^{(1)}) f(h^{(2)} l^{(2)}, m),$$

- (3) (Twisted module condition) for all $h, l \in H$ and $a \in A$ we have

$$\sum_{(h)(l)} (a^{l^{(1)}})^{h^{(1)}} f(h^{(2)}, l^{(2)}) = \sum_{(h)(l)} f(h^{(1)}, l^{(1)}) a^{h^{(2)} l^{(2)}}.$$

From now on we assume that H is the enveloping algebra $U(\mathfrak{g})$ of a Lie algebra $\mathfrak{g}$. In this case, item (1) of the definition of weak action implies that

$$(ab)^x = a^x b + ab^x$$

for each $x \in \mathfrak{g}$ and $a, b \in A$. So, a weak action determines a k -linear map

$$\delta: \mathfrak{g} \rightarrow \text{Der}_k(A)$$

by $\delta(x)(a) = a^x$. Moreover if $(h, a) \mapsto a^h$ is an action, then δ is a homomorphism of Lie algebras. Conversely, given a k -linear map $\delta: \mathfrak{g} \rightarrow \text{Der}_k(A)$, there exists a (generality non-unique) weak action of $U(\mathfrak{g})$ on A such that $\delta(x)(a) = a^x$. When δ is a homomorphism of Lie algebras, there is a unique action of $U(\mathfrak{g})$ on A such that $\delta(x)(a) = a^x$. For a proof of these facts see [B-C-M]. It is easy to see that each normal cocycle

$$f: U(\mathfrak{g}) \otimes_k U(\mathfrak{g}) \rightarrow A$$

is convolution invertible. For a proof see [G-G1, Remark 1.1].

Next we recall some results and notations from [G-G1] that we will need later. Let K be a subalgebra of A which is stable under the weak action of $\mathfrak{g}$ (that is $\lambda^x \in K$ for all $\lambda \in K$ and $x \in \mathfrak{g}$) and let $E = A \#_f U(\mathfrak{g})$ be a crossed product. We are going to modify the sign of some boundary maps in order to obtain simple expressions for the comparison maps.

To begin, we fix some notations:

- (1) The unadorned tensor product $\otimes$ means the tensor product $\otimes_K$ over K ,
- (2) For $B = A$ or $B = E$ and each $r \in \mathbb{N}$, we write $\overline{B} = B/K$,

$$B^r = B \otimes \cdots \otimes B \text{ (} r \text{ times)} \text{ and } \overline{B}^r = \overline{B} \otimes \cdots \otimes \overline{B} \text{ (} r \text{ times)}.$$

Moreover, for $b \in B$ we also let $\overline{b}$ denote the class of b in $\overline{B}$.

- (3) For each Lie algebra $\mathfrak{g}$ and $s \in \mathbb{N}$, we write $\mathfrak{g}^{\wedge s} = \mathfrak{g} \wedge \cdots \wedge \mathfrak{g}$ (s times).
- (4) Throughout this paper we will write $\mathbf{a}_{1r}$ for $a_1 \otimes \cdots \otimes a_r \in A^r$ and $\mathbf{x}_{1s}$ for $x_1 \wedge \cdots \wedge x_s \in \mathfrak{g}^{\wedge s}$
- (5) For $\mathbf{a}_{1r}$ and $0 \leq i < j \leq r$, we write $\mathbf{a}_{ij} = a_i \otimes \cdots \otimes a_j$.
- (6) For $\mathbf{x}_{1s}$ and $1 \leq i \leq s$, we write $\mathbf{x}_{1\widehat{i}s} = x_1 \wedge \cdots \wedge \widehat{x_i} \wedge \cdots \wedge x_s$.
- (7) For $\mathbf{x}_{1s}$ and $1 \leq i < j \leq s$, we write $\mathbf{x}_{1\widehat{i}\widehat{j}s} = x_1 \wedge \cdots \wedge \widehat{x_i} \wedge \cdots \wedge \widehat{x_j} \wedge \cdots \wedge x_s$.

Let Y_* be the graded algebra generated by A and the elements y_x, z_x ($x \in \mathfrak{g}$) in degree zero, the elements e_x ($x \in \mathfrak{g}$) in degree one, and the relations

$$\begin{aligned} y_{\lambda x + x'} &= \lambda y_x + y_{x'}, & y_x a &= a^x + a y_x, & e_{x'} y_x &= y_x e_{x'} + e_{[x', x]_{\mathfrak{g}}}, \\ z_{\lambda x + x'} &= \lambda z_x + z_{x'}, & z_x a &= a^x + a z_x, & e_{x'} z_x &= z_x e_{x'}, \\ e_{\lambda x + x'} &= \lambda e_x + e_{x'}, & e_x a &= a e_x, & e_x^2 &= 0, \\ y_{x'} y_x &= y_x y_{x'} + y_{[x', x]_{\mathfrak{g}}} + f(x', x) - f(x, x'), \\ z_{x'} y_x &= y_x z_{x'} + z_{[x', x]_{\mathfrak{g}}} + f(x', x) - f(x, x'), \\ z_{x'} z_x &= z_x z_{x'} + z_{[x', x]_{\mathfrak{g}}} + f(x', x) - f(x, x'), \end{aligned}$$

where $[x', x]_{\mathfrak{g}}$ denotes the Lie bracket of x' and x in $\mathfrak{g}$. Note that E is a subalgebra of Y_* via the embedding that takes $a \in A$ to a and $1 \# x$ to y_x for all $x \in \mathfrak{g}$. This gives rise to an structure of left E -module on Y_* . Similarly we consider Y_* as a right E -module via the embedding of E in Y_* that takes $a \in A$ to a and $1 \# x$ to z_x for all $x \in \mathfrak{g}$.

Let $(g_i)_{i \in I}$ be a basis of $\mathfrak{g}$ with indexes running on an ordered set I . For each $i \in I$ let us write $y_i = y_{g_i}$, $z_i = z_{g_i}$, $e_i = e_{g_i}$ and $\rho_i = z_i - y_i$.

Theorem 1.1. *Each Y_s is a free left E -module with basis*

$$\rho_{i_1}^{m_1} e_{i_1}^{\delta_1} \cdots \rho_{i_l}^{m_l} e_{i_l}^{\delta_l} \quad \left(\begin{array}{l} l \geq 0, i_1 < \cdots < i_l \in I, m_j \geq 0, \delta_j \in \{0, 1\} \\ m_j + \delta_j > 0, \delta_1 + \cdots + \delta_l = s \end{array} \right).$$

Proof. Let $\tilde{\mathfrak{g}}$ be the direct sum of two copies $\{y_x : x \in \mathfrak{g}\}$ and $\{z_x : x \in \mathfrak{g}\}$ of $\mathfrak{g}$, endowed with the bracket given by

$$[y_{x'}, y_x]_{\tilde{\mathfrak{g}}} = y_{[x', x]_{\mathfrak{g}}} \quad \text{and} \quad [z_{x'}, z_x]_{\tilde{\mathfrak{g}}} = [z_{x'}, y_x]_{\tilde{\mathfrak{g}}} = z_{[x', x]_{\mathfrak{g}}}.$$

Note that $\tilde{\mathfrak{g}}$ is the semi-direct sum arising from the adjoint action of $\mathfrak{g}$ on itself. Let $\pi: U(\tilde{\mathfrak{g}}) \rightarrow U(\mathfrak{g})$ be the algebra map defined by $\pi(y_x) = \pi(z_x) = x$. Let $\Lambda(\mathfrak{g})$ be the exterior algebra generated by $\mathfrak{g}$. That is, the algebra generated by the elements e_x , with $x \in \mathfrak{g}$, and the relations $e_{\lambda x + x'} = \lambda e_x + e_{x'}$ and $e_x^2 = 0$, with $\lambda \in k$ and $x, x' \in \mathfrak{g}$. Let us consider the action of $U(\tilde{\mathfrak{g}})$ on $\Lambda(\mathfrak{g})$ determined by $e_{x'}^{y_x} = e_{[x, x']_{\mathfrak{g}}}$ and $e_{x'}^{z_x} = 0$. The enveloping algebra $U(\tilde{\mathfrak{g}})$ of $\tilde{\mathfrak{g}}$ acts weakly on $A \otimes_k \Lambda(\mathfrak{g})$ via

$$(a \otimes_k e)^u = \sum_{(u)} a^{\pi(u^{(1)})} \otimes_k e^{u^{(2)}} \quad (a \in A, e \in \Lambda(\mathfrak{g}) \text{ and } u \in U(\tilde{\mathfrak{g}})).$$

Moreover, the map

$$\tilde{f}: U(\tilde{\mathfrak{g}}) \times U(\tilde{\mathfrak{g}}) \longrightarrow A \otimes_k \Lambda(\mathfrak{g}),$$

defined by $\tilde{f}(u, v) = f(\pi(u), \pi(v)) \otimes_k 1$, is a normal 2-cocycle which satisfies the twisted module condition. Let

$$\eta: Y'_* \longrightarrow (A \otimes_k \Lambda(\mathfrak{g})) \#_{\tilde{f}} U(\tilde{\mathfrak{g}})$$

be the homomorphism of algebras defined by $\eta(a) = (a \otimes_k 1) \# 1$ for all $a \in A$ and $\eta(y_x) = (1 \otimes_k 1) \# y_x$, $\eta(z_x) = (1 \otimes_k 1) \# z_x$ and $\eta(e_x) = (1 \otimes_k e_x) \# 1$ for all $x \in \mathfrak{g}$. Because of the Poincaré-Birkhoff-Witt theorem,

$$\eta(y_{j_1}^{n_1} \cdots y_{j_h}^{n_h} \rho_{i_1}^{m_1} \cdots \rho_{i_l}^{m_l}) \quad \left(\begin{array}{l} h, l \geq 0, j_1 < \cdots < j_h, \\ i_1 < \cdots < i_l \in I \text{ and } m_j, n_j \geq 0 \end{array} \right),$$

is a basis of $(A \otimes_k \Lambda(\mathfrak{g})) \#_{\tilde{f}} U(\tilde{\mathfrak{g}})$ as a left $A \otimes_k \Lambda(\mathfrak{g})$ -module. The theorem follows easily from this fact. $\square$

Remark 1.2. A similar argument shows that each Y_s is a free right E -module with the same basis.

Theorem 1.3. *Let $\tilde{\mu}: Y_0 \rightarrow E$ be the algebra map defined by $\tilde{\mu}(a) = a$ for $a \in A$ and $\tilde{\mu}(y_x) = \tilde{\mu}(z_x) = 1 \# x$ for $x \in \mathfrak{g}$. There is a unique derivation $\partial_*: Y_* \rightarrow Y_{*-1}$ such that $\partial(e_x) = z_x - y_x$ for $x \in \mathfrak{g}$. Moreover, the chain complex of E -bimodules*

$$E \xleftarrow{\tilde{\mu}} Y_0 \xleftarrow{\partial_1} Y_1 \xleftarrow{\partial_2} Y_2 \xleftarrow{\partial_3} Y_3 \xleftarrow{\partial_4} Y_4 \xleftarrow{\partial_5} Y_5 \xleftarrow{\partial_6} \cdots$$

is contractible as a complex of (E, A) -bimodules. A chain contracting homotopy

$$\sigma_0^{-1}: E \longrightarrow Y_0, \quad \sigma_{s+1}^{-1}: Y_s \longrightarrow Y_{s+1} \quad (s \geq 0),$$

is given by

$$\sigma^{-1}(1) = 1, \\ \sigma^{-1}(\rho_{i_1}^{m_1} e_{i_1}^{\delta_1} \cdots \rho_{i_l}^{m_l} e_{i_l}^{\delta_l}) = \begin{cases} \rho_{i_1}^{m_1-1} e_{i_1} \rho_{i_2}^{m_2} e_{i_2}^{\delta_2} \cdots \rho_{i_l}^{m_l} e_{i_l}^{\delta_l} & \text{if } \delta_1 = 0, \\ 0 & \text{if } \delta_1 = 1, \end{cases}$$

where we assume that $i_1 < \cdots < i_l$, $\delta_1 + \cdots + \delta_l = s$ and $m_l + \delta_l > 0$.

Proof. A direct computation shows that

$$- \tilde{\mu} \circ \sigma^{-1}(1) = \tilde{\mu}(1) = 1,$$

- $\sigma^{-1} \circ \tilde{\mu}(1) = \sigma^{-1}(1) = 1$ and $\partial \circ \sigma^{-1}(1) = \partial(0) = 0$,
- If $\mathbf{x} = \rho_{i_1}^{m_1} \mathbf{x}'$, where $m_1 > 0$ and $\mathbf{x}' = \rho_{i_2}^{m_2} \cdots \rho_{i_l}^{m_l}$ with $i_1 < \cdots < i_l$, then

$$\sigma^{-1} \circ \tilde{\mu}(\mathbf{x}) = \sigma^{-1}(0) = 0 \quad \text{and} \quad \partial \circ \sigma^{-1}(\mathbf{x}) = \partial(\rho_{i_1}^{m_1-1} e_{i_1} \mathbf{x}') = \mathbf{x},$$
- Let $\mathbf{x} = \rho_{i_1}^{m_1} e_{i_1}^{\delta_1} \mathbf{x}'$, where $m_l + \delta_l > 0$ and $\mathbf{x}' = \rho_{i_2}^{m_2} e_{i_2}^{\delta_2} \cdots \rho_{i_l}^{m_l} e_{i_l}^{\delta_l}$ with $i_1 < \cdots < i_l$ and $\delta_1 + \cdots + \delta_l = s > 0$. If $\delta_1 = 0$, then

$$\begin{aligned} \sigma^{-1} \circ \partial(\mathbf{x}) &= \sigma^{-1}(\rho_{i_1}^{m_1} \partial(\mathbf{x}')) = \rho_{i_1}^{m_1-1} e_{i_1} \partial(\mathbf{x}'), \\ \partial \circ \sigma^{-1}(\mathbf{x}) &= \partial(\rho_{i_1}^{m_1-1} e_{i_1} \mathbf{x}') = \mathbf{x} - \rho_{i_1}^{m_1-1} e_{i_1} \partial(\mathbf{x}'), \end{aligned}$$

and if $\delta_1 = 1$, then

$$\begin{aligned} \sigma^{-1} \circ \partial(\mathbf{x}) &= \sigma^{-1}(\rho_{i_1}^{m_1+1} \mathbf{x}' - \rho_{i_1}^{m_1} e_{i_1} \partial(\mathbf{x}')) = \mathbf{x}, \\ \partial \circ \sigma^{-1}(\mathbf{x}) &= \partial(0) = 0. \end{aligned}$$

The result follows immediately from all these facts. $\square$

For each $s \geq 0$ we consider $E \otimes_k \mathfrak{g}^{\wedge s}$ as a right K -module via $(\mathbf{c} \otimes_k \mathbf{x})\lambda = \mathbf{c}\lambda \otimes_k \mathbf{x}$. For $r, s \geq 0$, let $X_{rs} = (E \otimes_k \mathfrak{g}^{\wedge s}) \otimes \overline{A}^r \otimes E$. The groups X_{rs} are E -bimodules in an obvious way. Let us consider the diagram of E -bimodules and E -bimodule maps

$$\begin{array}{ccccccc} & & \vdots & & & & \\ & & \downarrow \partial_3 & & & & \\ Y_2 & \xleftarrow{\mu_2} & X_{02} & \xleftarrow{d_{12}^0} & X_{12} & \xleftarrow{d_{22}^0} & \cdots \\ & & \downarrow \partial_2 & & & & \\ Y_1 & \xleftarrow{\mu_1} & X_{01} & \xleftarrow{d_{11}^0} & X_{11} & \xleftarrow{d_{21}^0} & \cdots \\ & & \downarrow \partial_1 & & & & \\ Y_0 & \xleftarrow{\mu_0} & X_{00} & \xleftarrow{d_{10}^0} & X_{10} & \xleftarrow{d_{20}^0} & \cdots, \end{array}$$

where $\mu_* : X_{0*} \rightarrow Y_*$ and $d_{**}^0 : X_{**} \rightarrow X_{*-1,*}$, are defined by:

$$\begin{aligned} \mu(1 \otimes_k \mathbf{x}_{1s} \otimes 1) &= e_{x_1} \cdots e_{x_s}, \\ d^0(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1) &= (-1)^s a_1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{2r} \otimes 1 \\ &\quad + \sum_{i=1}^{r-1} (-1)^{i+s} \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1,i-1} \otimes a_i a_{i+1} \otimes \mathbf{a}_{i+1,r} \otimes 1 \\ &\quad + (-1)^{r+s} \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1,r-1} \otimes a_r, \end{aligned}$$

Each horizontal complex in this diagram is contractible as a complex of (E, K) -bimodules. A chain contracting homotopy is the family

$$\sigma_{0s}^0 : Y_s \longrightarrow X_{0s}, \quad \sigma_{r+1,s}^0 : X_{rs} \longrightarrow X_{r+1,s} \quad (r \geq 0),$$

of (E, K) -bimodule maps, defined by

$$\sigma^0(e_{x_1} \cdots e_{x_s} z_{x_{s+1}} \cdots z_{x_n}) = \sum_j a_j \otimes_k \mathbf{x}_{1s} \otimes 1 \# w_j,$$

where $\sum_j a_j \# w_j = (1 \# x_{s+1}) \cdots (1 \# x_n)$, and

$$\sigma^0(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes a_{r+1} \# w) = (-1)^{r+s+1} \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1,r+1} \otimes 1 \# w \quad (r \geq 0).$$

(In order to see that the σ^0 's are right K -linear it is necessary to use that K is stable under the action of $\mathfrak{g}$). Moreover, each X_{rs} is a projective E -bimodule relative to

the family of all epimorphism of E -bimodules which split as (E, K) -bimodule maps. We define E -bimodule maps

$$d_{rs}^l: X_{rs} \longrightarrow X_{r+l-1, s-l} \quad (r \geq 0 \text{ and } 1 \leq l \leq s),$$

recursively by:

$$d^l(\mathbf{y}) = \begin{cases} -\sigma^0 \circ \partial \circ \mu(\mathbf{y}) & \text{if } l = 1 \text{ and } r = 0, \\ -\sigma^0 \circ d^1 \circ d^0(\mathbf{y}) & \text{if } l = 1 \text{ and } r > 0, \\ -\sum_{j=1}^{l-1} \sigma^0 \circ d^{l-j} \circ d^j(\mathbf{y}) & \text{if } l > 1 \text{ and } r = 0, \\ -\sum_{j=0}^{l-1} \sigma^0 \circ d^{l-j} \circ d^j(\mathbf{y}) & \text{if } l > 1 \text{ and } r > 0, \end{cases}$$

where $\mathbf{y} = 1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1$.

Theorem 1.4. *The complex*

$$(1) \quad E \xleftarrow{\bar{\mu}} X_0 \xleftarrow{d_1} X_1 \xleftarrow{d_2} X_2 \xleftarrow{d_3} X_3 \xleftarrow{d_4} X_4 \xleftarrow{d_5} X_5 \xleftarrow{d_6} \cdots,$$

where

$$\bar{\mu}(1 \otimes 1) = 1, \quad X_n = \bigoplus_{r+s=n} X_{rs} \quad \text{and} \quad d_n = \sum_{\substack{r+s=n \\ r+l>0}} \sum_{l=0}^s d_{rs}^l,$$

is a projective resolution of the E -bimodule E , relative to the family of all epimorphism of E -bimodules which split as (E, K) -bimodule maps. Moreover an explicit contracting homotopy

$$\bar{\sigma}_0: E \longrightarrow X_0, \quad \bar{\sigma}_{n+1}: X_n \longrightarrow X_{n+1} \quad (n \geq 0),$$

of (1), as a complex of (E, K) -bimodules, is given by

$$\bar{\sigma}_0 = \sigma^0 \circ \sigma_0^{-1} \quad \text{and} \quad \bar{\sigma}_{n+1} = - \sum_{l=0}^{n+1} \sigma_{l, n-l+1}^l \circ \sigma_{n+1}^{-1} \circ \mu_n + \sum_{r=0}^n \sum_{l=0}^{n-r} \sigma_{r+l+1, n-l-r}^l,$$

where

$$\sigma_{l, s-l}^l: Y_s \rightarrow X_{l, s-l} \quad \text{and} \quad \sigma_{r+l+1, s-l}^l: X_{rs} \rightarrow X_{r+l+1, s-l} \quad (0 < l \leq s, r \geq 0)$$

are recursively defined by

$$\sigma^l = - \sum_{j=0}^{l-1} \sigma^0 \circ d^{l-j} \circ \sigma^j.$$

Proof. By [G-G2, Corollary A.2]. □

The boundary maps of the projective resolution of E that we just found are defined recursively. Next we give closed formulas for them.

Theorem 1.5. For $x_i, x_j \in \mathfrak{g}$, we put $\widehat{f}_{ij} = f(x_i, x_j) - f(x_j, x_i)$. We have:

$$\begin{aligned}
d^1(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1) &= \sum_{i=1}^s (-1)^{i+1} \#x_i \otimes_k \mathbf{x}_{1\widehat{i}s} \otimes \mathbf{a}_{1r} \otimes 1 \\
&\quad + \sum_{i=1}^s (-1)^i \otimes_k \mathbf{x}_{1\widehat{i}s} \otimes \mathbf{a}_{1r} \otimes 1 \#x_i \\
&\quad + \sum_{\substack{i=1 \\ 1 \leq h \leq r}}^s (-1)^i \otimes_k \mathbf{x}_{1\widehat{i}s} \otimes \mathbf{a}_{1, h-1} \otimes a_h^{x_i} \otimes \mathbf{a}_{h+1, r} \otimes 1 \\
&\quad + \sum_{1 \leq i < j \leq s} (-1)^{i+j} \otimes_k [x_i, x_j]_{\mathfrak{g}} \wedge \mathbf{x}_{1\widehat{ij}s} \otimes \mathbf{a}_{1r} \otimes 1, \\
d^2(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1) &= \sum_{\substack{1 \leq i < j \leq s \\ 0 \leq h \leq r}} (-1)^{i+j+h+s} \otimes_k \mathbf{x}_{1\widehat{ij}s} \otimes \mathbf{a}_{1h} \otimes \widehat{f}_{ij} \otimes \mathbf{a}_{h+1, r} \otimes 1
\end{aligned}$$

and $d^l = 0$ for all $l \geq 3$.

Proof. The proof of [G-G1, Theorem 3.3] works in our more general context. $\square$

2. THE COMPARISON MAPS

In this section we introduce and study comparison maps between (X_*, d_*) and the canonical normalized Hochschild resolution $(E \otimes \overline{E}^* \otimes E, b'_*)$ of the K -algebra E . It is well known that there are morphisms of E -bimodule complexes

$$\theta_* : (X_*, d_*) \longrightarrow (E \otimes \overline{E}^* \otimes E, b'_*) \quad \text{and} \quad \vartheta_* : (E \otimes \overline{E}^* \otimes E, b'_*) \longrightarrow (X_*, d_*),$$

such that $\theta_0 = \vartheta_0 = \text{id}_{E \otimes E}$ and that these morphisms are inverse one of each other up to homotopy. They can be recursively defined by $\theta_0 = \vartheta_0 = \text{id}_{E \otimes E}$ and

$$\theta(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1) = (-1)^n \theta \circ d(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1) \otimes 1$$

and

$$\vartheta(1 \otimes \mathbf{c}_{1n} \otimes 1) = \overline{\sigma} \circ \vartheta \circ b'(1 \otimes \mathbf{c}_{1n} \otimes 1),$$

for $n \geq 1$, where $r + s = n$ and $\mathbf{c}_{1n} = c_1 \otimes \cdots \otimes c_n \in \overline{E}^n$. The following result was established without proof in [G-G1].

Proposition 2.1. We have:

$$\theta(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1) = \sum_{\tau \in \mathfrak{S}_s} \text{sg}(\tau) \otimes (1 \# x_{\tau(1)} \otimes \cdots \otimes 1 \# x_{\tau(s)}) * \mathbf{a}_{1r} \otimes 1,$$

where $\mathfrak{S}_s$ is the symmetric group in s elements and $*$ denotes the shuffle product, which is defined by

$$(\beta_1 \otimes \cdots \otimes \beta_s) * (\beta_{s+1} \otimes \cdots \otimes \beta_n) = \sum_{\sigma \in \{(s, n-s)\text{-shuffles}\}} \text{sg}(\sigma) \beta_{\sigma(1)} \otimes \cdots \otimes \beta_{\sigma(n)}.$$

Proof. We proceed by induction on $n = r + s$. The case $n = 0$ is obvious. Suppose that $r + s = n$ and the result is valid for θ_{n-1} . By the recursive definition of θ ,

Theorem 1.5, and the inductive hypothesis,

$$\begin{aligned}
\theta(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1) &= (-1)^n \theta \circ d(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1) \otimes 1 \\
&= (-1)^n \theta \circ (d^0 + d^1 + d^2)(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1) \otimes 1 \\
&= \theta(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1,r-1} \otimes a_r) \otimes 1 \\
&\quad + \theta \left(\sum_{i=1}^s (-1)^{i+n} \otimes_k \mathbf{x}_{1\widehat{i}s} \otimes \mathbf{a}_{1r} \otimes 1 \# x_i \right) \otimes 1.
\end{aligned}$$

The desired result follows now using again the inductive hypothesis. $\square$

Lemma 2.2. *Let $(g_i)_{i \in I}$ be the basis of $\mathfrak{g}$ considered in Theorem 1.1. As in that theorem, let us write $e_i = e_{g_i}$ for each $i \in I$. The following facts hold:*

- (1) $\bar{\sigma}_{n+1} \circ \bar{\sigma}_n = 0$ for all $n \geq 0$,
- (2) $\sigma^l((E \otimes_k \mathfrak{g}^{\wedge s}) \otimes \bar{A}^r \otimes K \# U(\mathfrak{g})) = 0$ for all $0 \leq l \leq s$,
- (3) $\sigma^l(e_{i_1} \cdots e_{i_n}) = 0$ for all $0 < l \leq n$,
- (4) $\sigma^l((E \otimes_k \mathfrak{g}^{\wedge s}) \otimes \bar{A}^r \otimes A) = 0$ for all $0 < l \leq s$,
- (5) $\sigma^{-1} \circ \mu(A \otimes_k \mathfrak{g}^{\wedge n} \otimes A) = 0$,
- (6) Assume that $i_1 < \cdots < i_n$. Then,

$$\sigma^{-1} \circ \mu(1 \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_n} \otimes 1 \# g_{i_{n+1}}) = \begin{cases} (-1)^n e_{i_1} \cdots e_{i_{n+1}} & \text{if } i_n < i_{n+1}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. (1) An inductive argument shows that there exist maps (which are left E -linear and right K -linear)

$$\gamma_{rs}^l : X_{r+1,s} \rightarrow X_{r+l,s-l},$$

such that $\sigma_{r+l+1,s-l}^l = \sigma_{r+l+1,s-l}^0 \circ \gamma_{rs}^l \circ \sigma_{rs}^0$. Because of $\sigma^0 \circ \sigma^0 = 0$, this implies that $\sigma^{l'} \circ \sigma^l = 0$, for all $l, l' \geq 0$. Thus,

$$\bar{\sigma}_{n+1} \circ \bar{\sigma}_n = \sum_{l=0}^{n+1} \sigma^l \circ \sigma^{-1} \circ \mu \circ \sigma^0 \circ \sigma^{-1} \circ \mu = 0,$$

where the last equality holds because $\mu \circ \sigma^0 = \text{id}$ and $\sigma^{-1} \circ \sigma^{-1} = 0$.

(2) Since $\sigma^l = \sigma^0 \circ \gamma^l \circ \sigma^0$ for $l > 0$, we can assume that $l = 0$. In this case the assertion follows immediately from the definition of σ^0 .

(3) By the definition of σ^0 and Theorem 1.5,

$$\sigma^0 \circ d^1 \circ \sigma^0(e_{i_1} \cdots e_{i_n}) = \sigma^0 \circ d^1(1 \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_n} \otimes 1) = 0$$

and

$$\sigma^0 \circ d^2 \circ \sigma^0(e_{i_1} \cdots e_{i_n}) = \sigma^0 \circ d^2(1 \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_n} \otimes 1) = 0.$$

Item (3) follows now easily by induction on l , since, by the recursive definition of σ^l and Theorem 1.5,

$$\sigma^1 = -\sigma^0 \circ d^1 \circ \sigma^0 \quad \text{and} \quad \sigma^l = -\sigma^0 \circ d^1 \circ \sigma^{l-1} - \sigma^0 \circ d^2 \circ \sigma^{l-2} \quad \text{for } l \geq 2.$$

(4) It is similar to the proof of item (3).

(5) Since $e_i a = a e_i$ for all $i \in I$ and $a \in A$,

$$\sigma^{-1} \circ \mu(a \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_n} \otimes a') = \sigma^{-1}(a e_{i_1} \cdots e_{i_n} a') = \sigma^{-1}(a a' e_{i_1} \cdots e_{i_n}) = 0,$$

where the last equality follows from the definition of σ^{-1} .

(6) We have

$$\begin{aligned}\sigma^{-1}\mu(1 \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_n} \otimes 1 \# g_{i_{n+1}}) &= \sigma^{-1}(e_{i_1} \cdots e_{i_n} z_{i_{n+1}}) \\ &= \sigma^{-1}(e_{i_1} \cdots e_{i_n} (y_{i_{n+1}} + \rho_{i_{n+1}})) \\ &= \sigma^{-1}(y_{i_{n+1}} e_{i_1} \cdots e_{i_n}) \\ &\quad + \sigma^{-1}(e_{i_1} \cdots e_{i_n} \rho_{i_{n+1}}),\end{aligned}$$

where $z_{i_{n+1}}$, $y_{i_{n+1}}$ and $\rho_{i_{n+1}}$ are as in Theorem 1.1. So, in order to finish the proof it suffices to note that $\sigma^{-1}(y_{i_{n+1}} e_{i_1} \cdots e_{i_n}) = 0$ and

$$\sigma^{-1}(e_{i_1} \cdots e_{i_n} \rho_{i_{n+1}}) = \begin{cases} (-1)^n e_{i_1} \cdots e_{i_{n+1}} & \text{if } i_n < i_{n+1}, \\ 0 & \text{otherwise,} \end{cases}$$

which follows immediately from the fact that

$$e_{i_j} \rho_{i_{n+1}} = \rho_{i_{n+1}} e_{i_j} + e_{[x_{i_j}, x_{i_{n+1}}]_{\mathfrak{g}}} \quad \text{for all } j \text{ such that } i_j > i_{n+1},$$

and from the definition of σ^{-1} . $\square$

Theorem 2.3. *Let $(g_i)_{i \in I}$ be the basis of $\mathfrak{g}$ considered in Theorem 1.1. Assume that $\mathbf{c}_{1n} = c_1 \otimes \cdots \otimes c_n \in \overline{E}^n$ is a simple tensor with $c_j \in A \cup \{1 \# g_i : i \in I\}$ for all $j \in \{1, \dots, n\}$. If there exist $0 \leq s \leq n$ and $i_1 < \cdots < i_s$ in I , such that $c_j = 1 \# g_{i_j}$ for $1 \leq j \leq s$ and $c_j \in A$ for $s < j \leq n$, then*

$$\vartheta(1 \otimes \mathbf{c}_{1n} \otimes 1) = 1 \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_s} \otimes \mathbf{c}_{s+1,n} \otimes 1.$$

Otherwise, $\vartheta(1 \otimes \mathbf{c}_{1n} \otimes 1) = 0$.

Proof. For all $n \geq 0$ we define P_n by $\mathbf{c}_{1n} \in P_n$ if there is $i_1 < \cdots < i_s$ in I such that $c_j = 1 \# g_{i_j}$ for $j \leq s$ and $c_j \in A$ for $j > s$. We now proceed by induction on n . The case $n = 0$ is immediate. Assume that the result is valid for ϑ_n . By item (1) of Lemma 2.2 and the recursive definition of ϑ_n , we have

$$\bar{\sigma} \circ \vartheta(\mathbf{c}'_{0n} \otimes 1) = \bar{\sigma} \circ \bar{\sigma} \circ \vartheta(b'(\mathbf{c}'_{0n} \otimes 1)) = 0,$$

and so

$$\vartheta(1 \otimes \mathbf{c}_{1,n+1} \otimes 1) = (-1)^{n+1} \bar{\sigma} \circ \vartheta(1 \otimes \mathbf{c}_{1,n+1}).$$

Assume that $c_j \in A \cup \{1 \# g_i : i \in I\}$ for all $j \in \{1, \dots, n+1\}$. In order to finish the proof it suffices to show that

- If $\mathbf{c}_{1,n+1} \notin P_{n+1}$, then $\bar{\sigma} \circ \vartheta(1 \otimes \mathbf{c}_{1,n+1}) = 0$,
 - If $\mathbf{c}_{1,n+1} = 1 \# g_{i_1} \otimes \cdots \otimes 1 \# g_{i_s} \otimes \mathbf{a}_{s+1,n+1} \in P_{n+1}$, then
- $$\bar{\sigma} \circ \vartheta(1 \otimes \mathbf{c}_{1,n+1}) = (-1)^{n+1} \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_s} \otimes \mathbf{a}_{s+1,n+1} \otimes 1.$$

If $\mathbf{c}_{1n} \notin P_n$, then $\vartheta(1 \otimes \mathbf{c}_{1,n+1}) = 0$ by the inductive hypothesis. It remains to consider the case $\mathbf{c}_{1n} \in P_n$. We divide this into three subcases.

1) If $\mathbf{c}_{1n} = 1 \# g_{i_1} \otimes \cdots \otimes 1 \# g_{i_s} \otimes \mathbf{a}_{s+1,n}$ and $c_{n+1} = a_{n+1} \in A$, then

$$\begin{aligned}\bar{\sigma} \circ \vartheta(1 \otimes \mathbf{c}_{1,n+1}) &= \bar{\sigma}(1 \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_s} \otimes \mathbf{a}_{s+1,n+1}) \\ &= \sigma^0(1 \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_s} \otimes \mathbf{a}_{s+1,n+1}) \\ &= (-1)^{n+1} \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_s} \otimes \mathbf{a}_{s+1,n+1} \otimes 1,\end{aligned}$$

by the inductive hypothesis, items (4) and (5) of Lemma 2.2, and the definitions of $\bar{\sigma}$ and σ^0 .

2) If $\mathbf{c}_{1n} = 1 \# g_{i_1} \otimes \cdots \otimes 1 \# g_{i_s} \otimes \mathbf{a}_{s+1,n}$ with $s < n$ and $c_{n+1} = 1 \# g_{i_{n+1}}$, then

$$\bar{\sigma} \circ \vartheta(1 \otimes \mathbf{c}_{1,n+1}) = \bar{\sigma}(1 \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_s} \otimes \mathbf{a}_{s+1,n} \otimes 1 \# g_{i_{n+1}}) = 0,$$

by the inductive hypothesis, the definition of $\bar{\sigma}$ and item (2) of Lemma 2.2.

3) If $\mathbf{c}_{1n} = 1 \# g_{i_1} \otimes \cdots \otimes 1 \# g_{i_n}$ and $c_{n+1} = 1 \# g_{i_{n+1}}$, then

$$\begin{aligned} \bar{\sigma} \circ \vartheta(1 \otimes \mathbf{c}_{1,n+1}) &= \bar{\sigma}(1 \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_n} \otimes 1 \# g_{i_{n+1}}) \\ &= -\sigma^0 \circ \sigma^{-1} \circ \mu(1 \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_n} \otimes 1 \# g_{i_{n+1}}) \\ &= \begin{cases} (-1)^{n+1} \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_{n+1}} \otimes 1 & \text{if } \mathbf{c}_{1,n+1} \in P_{n+1}, \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

by the inductive hypothesis, items (2), (3) and (6) of Lemma 2.2, and the definitions of $\bar{\sigma}$ and σ^0 . $\square$

3. THE HOCHSCHILD COHOMOLOGY

Let $E = A \#_f U(\mathfrak{g})$ and M an E -bimodule. In this section we obtain a cochain complex $(\bar{X}_K^*(M), \bar{d}^*)$, simpler than the canonical one, giving the Hochschild cohomology of the K -algebra E with coefficients in M . When $K = k$ our result reduce to the one obtained in [G-G1, Section 5]. Then, we obtain an expression that gives the cup product of the Hochschild cohomology of E in terms of $(\bar{X}_K^*(E), \bar{d}^*)$. As usual, given $c \in E$ and $m \in M$, we let $[m, c]$ denote the commutator $mc - cm$.

3.1. **The complex $(\bar{X}_K^*(M), \bar{d}^*)$.** For $r, s \geq 0$, let

$$\bar{X}_K^{rs}(M) = \text{Hom}_{K^e}(\bar{A}^r \otimes_k \mathfrak{g}^{\wedge s}, M),$$

where $\bar{A}^r \otimes_k \mathfrak{g}^{\wedge s}$ is considered as a K -bimodule via the canonical actions on $\bar{A}^r$. We define the morphism

$$\bar{d}_l^{rs} : \bar{X}_K^{r+l-1, s-l}(M) \longrightarrow \bar{X}_K^{rs}(M) \quad (\text{with } 0 \leq l \leq \min(2, s) \text{ and } r+l > 0),$$

by:

$$\begin{aligned} \bar{d}_0(\varphi)(\mathbf{a}_{1r} \otimes_k \mathbf{x}_{1s}) &= a_1 \varphi(\mathbf{a}_{2r} \otimes_k \mathbf{x}_{1s}) \\ &\quad + \sum_{i=1}^{r-1} (-1)^i \varphi(\mathbf{a}_{1,i-1} \otimes a_i a_{i+1} \otimes \mathbf{a}_{i+2,r} \otimes_k \mathbf{x}_{1s}) \\ &\quad + (-1)^r \varphi(\mathbf{a}_{1,r-1} \otimes_k \mathbf{x}_{1s}) a_r, \\ \bar{d}_1(\varphi)(\mathbf{a}_{1r} \otimes_k \mathbf{x}_{1s}) &= \sum_{i=1}^s (-1)^{i+r} [\varphi(\mathbf{a}_{1r} \otimes_k \mathbf{x}_{1\widehat{i}s}), 1 \# x_i] \\ &\quad + \sum_{\substack{i=1 \\ 1 \leq h \leq r}}^s (-1)^{i+r} \varphi(\mathbf{a}_{1,h-1} \otimes a_h^{x_i} \otimes \mathbf{a}_{h+1,r} \otimes_k \mathbf{x}_{1\widehat{i}s}) \\ &\quad + \sum_{1 \leq i < j \leq s} (-1)^{i+j+r} \varphi(\mathbf{a}_{1r} \otimes_k [x_i, x_j]_{\mathfrak{g}} \wedge \mathbf{x}_{1\widehat{i}\widehat{j}s}) \end{aligned}$$

and

$$\bar{d}_2(\varphi)(\mathbf{a}_{1r} \otimes_k \mathbf{x}_{1s}) = \sum_{\substack{1 \leq i < j \leq s \\ 0 \leq h \leq r}} (-1)^{i+j+h} \varphi(\mathbf{a}_{1h} \otimes \widehat{f}_{ij} \otimes \mathbf{a}_{h+1,r} \otimes_k \mathbf{x}_{1\widehat{i}\widehat{j}s}),$$

where $\widehat{f}_{ij} = f(x_i, x_j) - f(x_j, x_i)$. Recall that $X_{rs} = (E \otimes_k \mathfrak{g}^{\wedge s}) \otimes \bar{A}^r \otimes E$. Applying the functor $\text{Hom}_{E^e}(-, M)$ to the complex (X_*, d_*) of Theorem 1.4, and using Theorem 1.5 and the identifications $\gamma^{rs} : \bar{X}_K^{rs}(M) \rightarrow \text{Hom}_{E^e}(X_{rs}, M)$, given by

$$\gamma(\varphi)(1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1) = (-1)^{rs} \varphi(\mathbf{a}_{1r} \otimes_k \mathbf{x}_{1s}),$$

we obtain the complex

$$\overline{X}_K^0(M) \xrightarrow{\overline{d}^1} \overline{X}_K^1(M) \xrightarrow{\overline{d}^2} \overline{X}_K^2(M) \xrightarrow{\overline{d}^3} \overline{X}_K^3(M) \xrightarrow{\overline{d}^4} \overline{X}_K^4(M) \xrightarrow{\overline{d}^5} \cdots,$$

where

$$\overline{X}_K^n(M) = \bigoplus_{r+s=n} \overline{X}_K^{rs}(M) \quad \text{and} \quad \overline{d}^n = \sum_{\substack{r+s=n \\ r+l>0}} \sum_{l=0}^{\min(s,2)} \overline{d}_l^{rs}.$$

Note that if $f(\mathfrak{g} \otimes_k \mathfrak{g}) \subseteq K$, then $(\overline{X}_K^*(M), \overline{d}^*)$ is the total complex of the double complex $(\overline{X}_K^{**}(M), \overline{d}_0^{**}, \overline{d}_1^{**})$.

Theorem 3.1. *The Hochschild cohomology $H_K^*(E, M)$, of the K -algebra E with coefficients in M , is the cohomology of $(\overline{X}_K^*(M), \overline{d}^*)$.*

Proof. It is an immediate consequence of the above discussion. $\square$

3.2. The comparison maps. The maps θ_* and ϑ_* , introduced in Section 2, induce quasi-isomorphisms

$$\overline{\theta}^* : (\text{Hom}_{K^e}(\overline{E}^*, M), b^*) \longrightarrow (\overline{X}_K^*(M), \overline{d}^*)$$

and

$$\overline{\vartheta}^* : (\overline{X}_K^*(M), \overline{d}^*) \longrightarrow (\text{Hom}_{K^e}(\overline{E}^*, M), b^*)$$

which are inverse one of each other up to homotopy.

Proposition 3.2. *We have*

$$\overline{\theta}(\psi)(\mathbf{a}_{1r} \otimes_k \mathbf{x}_{1s}) = \sum_{\tau \in \mathfrak{S}_s} (-1)^{rs} \text{sg}(\tau) \psi \left((1 \# x_{\tau(1)} \otimes \cdots \otimes 1 \# x_{\tau(s)}) * \mathbf{a}_{1r} \right)$$

Proof. This follows immediately from Proposition 2.1. $\square$

In the sequel we consider that $\overline{X}_K^{rs} \subseteq \overline{X}_K^{r+s}$ in the canonical way.

Theorem 3.3. *Let $(g_i)_{i \in I}$ be the basis of $\mathfrak{g}$ considered in Theorem 1.1 and let $\varphi \in \overline{X}_K^{rs}$. Assume that $\mathbf{c}_{1,r+s} = c_1 \otimes \cdots \otimes c_{r+s} \in \overline{E}^{r+s}$ is a simple tensor with $c_j \in A \cup \{1 \# g_i : i \in I\}$ for all $j \in \{1, \dots, r+s\}$. If $c_j = 1 \# g_{i_j}$ with $i_1 < \cdots < i_s$ in I for $1 \leq j \leq s$ and $c_j \in A$ for $s < j \leq r+s$, then*

$$\overline{\vartheta}(\varphi)(\mathbf{c}_{1,r+s}) = (-1)^{rs} \varphi(\mathbf{c}_{s+1,r+s} \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_s}).$$

Otherwise, $\overline{\vartheta}(\varphi)(\mathbf{c}_{1,r+s}) = 0$.

Proof. This follows immediately from Theorem 2.3. $\square$

As usual, in the following subsection we will write $\text{HH}_K^*(E)$ instead of $H_K^*(E, E)$.

3.3. The cup product. Recall that the cup product of $\text{HH}_K^*(E)$ is given in terms of $(\text{Hom}_{K^e}(\overline{E}^*, E), b^*)$, by

$$(\psi \smile \psi')(\mathbf{c}_{1,m+n}) = \psi(\mathbf{c}_{1m}) \psi'(\mathbf{c}_{m+1,m+n}),$$

where $\psi \in \text{Hom}_{K^e}(\overline{E}^m, E)$ and $\psi' \in \text{Hom}_{K^e}(\overline{E}^n, E)$. In this subsection we compute the cup product in terms of the small complex $(\overline{X}_K^*(E), \overline{d}^*)$. Given $\varphi \in \overline{X}_K^{rs}(E)$ and $\varphi' \in \overline{X}_K^{r's'}(E)$ we define $\varphi \bullet \varphi' \in \overline{X}_K^{r+r',s+s'}(E)$ by

$$(\varphi \bullet \varphi')(\mathbf{a}_{1r''} \otimes_k \mathbf{x}_{1s''}) = \sum_{1 \leq j_1 < \cdots < j_s \leq s''} \text{sg}(j_{1s}) \varphi(\mathbf{a}_{1r} \otimes_k \mathbf{x}_{j_{1s}}) \varphi'(\mathbf{a}_{r+1,r''} \otimes_k \mathbf{x}_{l_{1s'}}),$$

where

- $\text{sg}(j_{1s}) = (-1)^{r's + \sum_{u=1}^s (j_u - u)}$,
- $r'' = r + r'$ and $s'' = s + s'$,
- $1 \leq l_1 < \dots < l_{s'} \leq s''$ denote the set defined by
$$\{j_1, \dots, j_s\} \cup \{l_1, \dots, l_{s'}\} = \{1, \dots, s''\},$$
- $\mathbf{x}_{j_{1s}} = x_{j_1} \wedge \dots \wedge x_{j_s}$ and $\mathbf{x}_{l_{1s'}} = x_{l_1} \wedge \dots \wedge x_{l_{s'}}$.

Theorem 3.4. *The cup product of $\text{HH}_K^*(E)$ is induced by the operation $\bullet$ in the complex $(\overline{X}_K^*(E), \overline{d}^*)$.*

Proof. Let $\varphi \in \overline{X}_K^{r,s}(E)$ and $\varphi' \in \overline{X}_K^{r',s'}(E)$. Let r'' and s'' be natural numbers satisfying $r'' + s'' = r + r' + s + s'$ and let $\mathbf{a}_{1r''} \otimes_k \mathbf{x}_{1s''} \in X_{r'',s''}^K$. Let $(g_i)_{i \in I}$ be the basis of $\mathfrak{g}$ considered in Theorem 1.1. Clearly we can assume that there exist $i_1 < \dots < i_{s''}$ in I such that $x_j = g_{i_j}$ for all $1 \leq j \leq s''$. By Proposition 3.2,

$$\overline{\theta}(\overline{\theta}(\varphi) \smile \overline{\theta}(\varphi'))(\mathbf{a}_{1r''} \otimes_k \mathbf{x}_{1s''}) = (\overline{\theta}(\varphi) \smile \overline{\theta}(\varphi'))(T)$$

where

$$T = \sum_{\tau \in \mathfrak{S}_{s''}} (-1)^{r''s''} \text{sg}(\tau) ((1 \# x_{\tau(1)}) \otimes \dots \otimes (1 \# x_{\tau(s'')})) * \mathbf{a}_{1r''}.$$

In order to finish the proof it suffices to note that by Theorem 3.3, this is zero if $r'' \neq r + r'$ and this is $(\varphi \bullet \varphi')(\mathbf{a}_{1r''} \otimes_k \mathbf{x}_{1s''})$ if $r'' = r + r'$. $\square$

4. THE HOCHSCHILD HOMOLOGY

Let $E = A \#_f U(\mathfrak{g})$ and M an E -bimodule. In this section we obtain a chain complex $(\overline{X}_*^K(M), \overline{d}_*)$, simpler than the canonical one, giving the Hochschild homology of the K -algebra E with coefficients in M . When $K = k$ our result reduce to the one obtained in [G-G1, Section 4]. Then, we obtain an expression that gives the cap product of $H_*^K(E, M)$ in terms of $(\overline{X}_K^*(E), \overline{d}^*)$ and $(\overline{X}_*^K(E, M), \overline{d}_*)$. As in the previous section $[m, c]$ denotes the commutator $mc - cm$ of $m \in M$ and $c \in E$.

4.1. The complex $(\overline{X}_*^K(M), \overline{d}_*)$. For $r, s \geq 0$, let

$$\overline{X}_{rs}^K(M) = \frac{M \otimes \overline{A}^r}{[M \otimes \overline{A}^r, K]} \otimes \mathfrak{g}^{\wedge s},$$

where $[M \otimes \overline{A}^r, K]$ is the k -vector space generated by the commutators $[m \otimes \mathbf{a}_{1r}, \lambda]$, with $\lambda \in K$ and $m \otimes \mathbf{a}_{1r} \in M \otimes \overline{A}^r$. We define the morphism

$$\overline{d}_{rs}^l: \overline{X}_{rs}^K(M) \longrightarrow \overline{X}_{r+l-1, s-l}^K(M) \quad (\text{with } 0 \leq l \leq \min(2, s) \text{ and } r + l > 0),$$

by:

$$\begin{aligned} \overline{d}_{rs}^0(\overline{m \otimes \mathbf{a}_{1r} \otimes_k \mathbf{x}_{1s}}) &= \overline{ma_1 \otimes \mathbf{a}_{2r} \otimes_k \mathbf{x}_{1s}} \\ &\quad + \sum_{i=1}^{r-1} (-1)^i \overline{m \otimes \mathbf{a}_{1,i-1} \otimes a_i a_{i+1} \otimes \mathbf{a}_{i+2,r} \otimes_k \mathbf{x}_{1s}} \\ &\quad + (-1)^r \overline{a_r m \otimes \mathbf{a}_{1,r-1} \otimes_k \mathbf{x}_{1s}}, \\ \overline{d}_{rs}^1(\overline{m \otimes \mathbf{a}_{1r} \otimes_k \mathbf{x}_{1s}}) &= \sum_{i=1}^s (-1)^{i+r} \overline{[(1 \# x_i), m] \otimes \mathbf{a}_{1r} \otimes_k \mathbf{x}_{1\widehat{i}s}} \\ &\quad + \sum_{\substack{i=1 \\ 1 \leq h \leq r}}^s (-1)^{i+r} \overline{m \otimes \mathbf{a}_{1,h-1} \otimes a_h^{x_i} \otimes \mathbf{a}_{h+1,r} \otimes_k \mathbf{x}_{1\widehat{i}s}} \end{aligned}$$

$$+ \sum_{1 \leq i < j \leq s} (-1)^{i+j+r} \overline{m \otimes \mathbf{a}_{1r}} \otimes_k [x_i, x_j]_{\mathfrak{g}} \wedge \mathbf{x}_{1\widehat{i}j_s}$$

and

$$\overline{d}^2(\overline{m \otimes \mathbf{a}_{1r}} \otimes_k \mathbf{x}_{1s}) = \sum_{\substack{1 \leq i < j \leq s \\ 0 \leq h \leq r}} (-1)^{i+j+h} \overline{m \otimes \mathbf{a}_{1h} \otimes \widehat{f_{ij}} \otimes \mathbf{a}_{h+1,r}} \otimes_k \mathbf{x}_{1\widehat{i}j_s},$$

where $\widehat{f_{ij}} = f(x_i, x_j) - f(x_j, x_i)$ and $\overline{m \otimes \mathbf{a}_{1r}}$ denotes the class of $m \otimes \mathbf{a}_{1r}$ in $M \otimes \overline{A^r} / [M \otimes \overline{A^r}, K]$, etcetera. Recall that $X_{rs} = (E \otimes_k \mathfrak{g}^{\wedge s}) \otimes \overline{A^r} \otimes E$ and let E^e be enveloping algebra of E . By tensoring on the left X_{rs} over E^e with M , and using Theorem 1.5 and the identifications $\gamma_{rs}: \overline{X}_{rs}^K(M) \rightarrow M \otimes_{E^e} X_{rs}$, given by

$$\gamma(\overline{m \otimes \mathbf{a}_{1r}} \otimes_k \mathbf{x}_{1s}) = (-1)^{rs} m \otimes_{E^e} (1 \otimes_k \mathbf{x}_{1s} \otimes \mathbf{a}_{1r} \otimes 1),$$

we obtain the complex

$$\overline{X}_0^K(M) \xleftarrow{\overline{d}_1} \overline{X}_1^K(M) \xleftarrow{\overline{d}_2} \overline{X}_2^K(M) \xleftarrow{\overline{d}_3} \overline{X}_3^K(M) \xleftarrow{\overline{d}_4} \overline{X}_4^K(M) \xleftarrow{\overline{d}_5} \cdots,$$

where

$$\overline{X}_n^K(M) = \bigoplus_{r+s=n} \overline{X}_{rs}^K(M) \quad \text{and} \quad \overline{d}_n = \sum_{\substack{r+s=n \\ r+l>0}} \sum_{l=0}^{\min(s,2)} \overline{d}_{rs}^l.$$

Note that if $f(\mathfrak{g} \otimes_k \mathfrak{g}) \subseteq K$, then $(\overline{X}_*^K(M), \overline{d}_*)$ is the total complex of the double complex $(\overline{X}_{**}^K(M), \overline{d}_{**}^0, \overline{d}_{**}^1)$.

Theorem 4.1. *The Hochschild homology $H_*^K(E, M)$, of the K -algebra E with coefficients in M , is the homology of $(\overline{X}_*^K(M), \overline{d}_*)$.*

Proof. It is an immediate consequence of the above discussion. $\square$

4.2. The comparison maps. The maps θ_* and ϑ_* , introduced in Section 2, induce quasi-isomorphisms

$$\overline{\theta}_*: (\overline{X}_*^K(M), \overline{d}_*) \longrightarrow \left(\frac{M \otimes \overline{E}^*}{[M \otimes \overline{E}^*, K]}, b_* \right)$$

and

$$\overline{\vartheta}_*: \left(\frac{M \otimes \overline{E}^*}{[M \otimes \overline{E}^*, K]}, b_* \right) \longrightarrow (\overline{X}_*^K(M), \overline{d}_*)$$

which are inverse one of each other up to homotopy.

Proposition 4.2. *We have*

$$\overline{\theta}(\overline{m \otimes \mathbf{a}_{1r}} \otimes_k \mathbf{x}_{1s}) = \sum_{\tau \in \mathfrak{S}_s} (-1)^{rs} \text{sg}(\tau) \overline{m \otimes (1 \# x_{\tau(1)} \otimes \cdots \otimes 1 \# x_{\tau(s)})} * \mathbf{a}_{1r}$$

Proof. This follows immediately from Proposition 2.1. $\square$

Theorem 4.3. *Let $(g_i)_{i \in I}$ be the basis of $\mathfrak{g}$ considered in Theorem 1.1. Assume that $\mathbf{c}_{1n} = c_1 \otimes \cdots \otimes c_n \in \overline{E}^n$ is a simple tensor with $c_j \in A \cup \{1 \# g_i : i \in I\}$ for all $j \in \{1, \dots, n\}$. If there exist $0 \leq s \leq n$ and $i_1 < \cdots < i_s$ in I , such that $c_j = 1 \# g_{i_j}$ for $1 \leq j \leq s$ and $c_j \in A$ for $s < j \leq n$, then*

$$\overline{\vartheta}(\overline{m \otimes \mathbf{c}_{1n}}) = (-1)^{s(n-s)} \overline{m \otimes \mathbf{c}_{s+1,n}} \otimes_k g_{i_1} \wedge \cdots \wedge g_{i_s}.$$

Otherwise, $\vartheta(\overline{m \otimes \mathbf{c}_{1n}}) = 0$.

Proof. This follows immediately from Theorem 2.3. $\square$

4.3. The cap product. Recall that the cap product

$$\mathrm{H}_p^K(E, M) \times \mathrm{HH}_K^q(E) \rightarrow \mathrm{H}_{p-q}^K(E, M) \quad (q \leq p),$$

is defined in terms of $\left(\frac{M \otimes \overline{E}^*}{[M \otimes \overline{E}^*, K]}, b_*\right)$ and $(\mathrm{Hom}_{K^e}(\overline{E}^*, E), b^*)$, by

$$\overline{m \otimes \mathbf{c}_{1p}} \frown \psi = \overline{m\psi(\mathbf{c}_{1q}) \otimes \mathbf{c}_{q+1,p}},$$

where $\psi \in \mathrm{Hom}_{K^e}(\overline{E}^q, E)$. In this subsection we compute the cup product in terms of the small complexes $(\overline{X}_*^K(M), \overline{d}_*)$ and $(\overline{X}_K^*(E), \overline{d}^*)$. Given

$$\overline{m \otimes \mathbf{a}_{1r}} \otimes_k \mathbf{x}_{1s} \in \overline{X}_{rs}^K(M) \quad \text{and} \quad \varphi' \in \overline{X}_{K'}^{r's'}(E) \quad \text{with } r \geq r' \text{ and } s \geq s',$$

we define $(\overline{m \otimes \mathbf{a}_{1r}} \otimes_k \mathbf{x}_{1s}) \bullet \varphi' \in \overline{X}_{r-r', s-s'}^K(M)$ by

$$(\overline{m \otimes \mathbf{a}_{1r}} \otimes_k \mathbf{x}_{1s}) \bullet \varphi' = \sum_{1 \leq j_1 < \dots < j_{s'} \leq s} \mathrm{sg}(j_{1s'}) \overline{m\varphi'(\mathbf{a}_{1r'} \otimes_k \mathbf{x}_{j_{1s'}})} \otimes \mathbf{a}_{r'+1, r} \otimes_k \mathbf{x}_{l_{1, s-s'}}.$$

where

- $\mathrm{sg}(j_{1s'}) = (-1)^{rs' + r's' + \sum_{u=1}^{s'} (j_u - u)}$,
- $1 \leq l_1 < \dots < l_{s-s'} \leq s$ denote the set defined by
$$\{j_1, \dots, j_{s'}\} \cup \{l_1, \dots, l_{s-s'}\} = \{1, \dots, s\},$$
- $\mathbf{x}_{j_{1s'}} = x_{j_1} \wedge \dots \wedge x_{j_{s'}}$ and $\mathbf{x}_{l_{1, s-s'}} = x_{l_1} \wedge \dots \wedge x_{l_{s-s'}}$.

Theorem 4.4. *The cap product*

$$\mathrm{H}_p^K(E, M) \times \mathrm{HH}_K^q(E) \rightarrow \mathrm{H}_{p-q}^K(E, M),$$

is induced by $\bullet$, in terms of the complexes $(\overline{X}_*^K(M), \overline{d}_*)$ and $(\overline{X}_K^*(E), \overline{d}^*)$.

Proof. Let $\overline{m \otimes \mathbf{a}_{1r}} \otimes_k \mathbf{x}_{1s} \in \overline{X}_{rs}^K(M)$ and $\varphi' \in \overline{X}_{K'}^{r's'}(E)$. Let $(g_i)_{i \in I}$ be the basis of $\mathfrak{g}$ considered in Theorem 1.1. Clearly we can assume that there exist $i_1 < \dots < i_s$ in I such that $x_j = g_{i_j}$ for all $1 \leq j \leq s$. By Proposition 4.2,

$$\overline{\vartheta}(\overline{\theta}(\overline{m \otimes \mathbf{a}_{1r}} \otimes_k \mathbf{x}_{1s}) \frown \overline{\vartheta}(\varphi')) = \overline{\vartheta}(T \frown \overline{\vartheta}(\varphi')),$$

where

$$T = \sum_{\sigma \in \mathfrak{S}_s} (-1)^{rs} \mathrm{sg}(\sigma) ((1 \# x_{\sigma(1)}) \otimes \dots \otimes (1 \# x_{\sigma(s)})) * \mathbf{a}_{1r}.$$

Hence, by Theorem 3.3, if $r' > r$ or $s' > s$, then

$$\overline{\vartheta}(\overline{\theta}(\overline{m \otimes \mathbf{a}_{1r}} \otimes_k \mathbf{x}_{1s}) \frown \overline{\vartheta}(\varphi')) = 0,$$

and, if $r' \leq r$ and $s' \leq s$, then

$$\overline{\vartheta}(\overline{\theta}(\overline{m \otimes \mathbf{a}_{1r}} \otimes_k \mathbf{x}_{1s}) \frown \overline{\vartheta}(\varphi')) = \sum_{1 \leq j_1 < \dots < j_{s'} \leq s} \overline{\vartheta}(m\varphi'(\mathbf{a}_{1r'} \otimes_k \mathbf{x}_{j_{1s'}}) \otimes T'_{l_{1, s-s'}}),$$

where

$$T'_{l_{1, s-s'}} = \sum_{\tau \in \mathfrak{S}_{s-s'}} (-1)^{rs+r's} \mathrm{sg}(\tau) ((1 \# x_{l_{\tau(1)}}) \otimes \dots \otimes (1 \# x_{l_{\tau(s-s')}})) * \mathbf{a}_{r'+1, r}.$$

In order to finish the proof it suffices to apply Theorem 4.3. $\square$

5. THE (CO)HOMOLOGY OF $S(V)\#_f U(\mathfrak{g})$

In this section we obtain a complexes $(\overline{Z}_*(M), \overline{\delta}_*)$ and $(\overline{Z}^*(M), \overline{\delta}^*)$, simpler than $(\overline{X}_K^*(M), \overline{d}^*)$ and $(\overline{X}_*^K(M), \overline{d}_*)$ respectively, giving the Hochschild homology of the K -algebra $E := A\#_f U(\mathfrak{g})$ with coefficients in an E -bimodule M

- $K = k$ and A is a symmetric algebra $S(V)$,
- $v^x \in k \oplus V$ for all $v \in V$ and $x \in \mathfrak{g}$,
- $f(x_1, x_2) \in k \oplus V$ for all $x_1, x_2 \in \mathfrak{g}$.

Then, we obtain an expression that gives the cup product of $H_K^*(E, M)$ in terms of $(\overline{Z}^*(E), \overline{\delta}^*)$, and we obtain an expression that gives the cap product of $H_*^K(E, M)$ in terms of $(\overline{Z}_*(M), \overline{\delta}_*)$ and $(\overline{Z}^*(E), \overline{\delta}^*)$.

For $r, s \geq 0$, let $Z_{rs} = E \otimes \mathfrak{g}^{\wedge s} \otimes V^{\wedge r} \otimes E$. The groups Z_{rs} are E -bimodules in an obvious way. Let

$$\delta_{rs}^l: Z_{rs} \longrightarrow Z_{r+l-1, s-l} \quad (0 \leq l \leq s \text{ and } r+l > 0),$$

be the E -bimodule morphisms defined by

$$\begin{aligned} \delta^0(1 \otimes \mathbf{x}_{1s} \otimes \mathbf{v}_{1r} \otimes 1) &= \sum_{i=1}^r (-1)^{i+s} (v_i \otimes \mathbf{x}_{1s} \otimes \mathbf{v}_{1\widehat{r}} \otimes 1 - 1 \otimes \mathbf{x}_{1s} \otimes \mathbf{v}_{1\widehat{r}} \otimes v_i), \\ \delta^1(1 \otimes \mathbf{x}_{1s} \otimes \mathbf{v}_{1r} \otimes 1) &= \sum_{i=1}^s (-1)^{i+1} \#x_i \otimes \mathbf{x}_{1\widehat{s}} \otimes \mathbf{v}_{1r} \otimes 1 \\ &\quad + \sum_{i=1}^s (-1)^i \otimes \mathbf{x}_{1\widehat{s}} \otimes \mathbf{v}_{1r} \otimes 1 \#x_i \\ &\quad + \sum_{\substack{i=1 \\ 1 \leq h \leq r}}^s (-1)^i \otimes \mathbf{x}_{1\widehat{s}} \otimes \mathbf{v}_{1, h-1} \wedge v_h^{\overline{x}_i} \wedge \mathbf{v}_{h+1, r} \otimes 1 \\ &\quad + \sum_{1 \leq i < j \leq s} (-1)^{i+j} \otimes [x_i, x_j]_{\mathfrak{g}} \wedge \mathbf{x}_{1\widehat{ij}} \otimes \mathbf{v}_{1r} \otimes 1 \end{aligned}$$

and

$$\delta^2(1 \otimes \mathbf{x}_{1s} \otimes \mathbf{v}_{1r} \otimes 1) = \sum_{1 \leq i < j \leq s} (-1)^{i+j+s} \otimes \mathbf{x}_{1\widehat{ij}} \otimes \widehat{f}_{ij} \wedge \mathbf{v}_{1r} \otimes 1,$$

where $\mathbf{v}_{hl} = v_h \wedge \cdots \wedge v_l$, $v_h^{\overline{x}_i}$ is the V -component of $v_h^{x_i}$ (that is $v_h^{\overline{x}_i} \in V$ and $v_h^{x_i} - v_h^{\overline{x}_i} \in k$) and $\widehat{f}_{ij} = f_V(x_i, x_j) - f_V(x_j, x_i)$ in which $f_V(x_j, x_i)$ is the V -component of $f(x_j, x_i)$.

Theorem 5.1. *The complex*

$$E \xleftarrow{\overline{\mu}} Z_0 \xleftarrow{\delta_1} Z_1 \xleftarrow{\delta_2} Z_2 \xleftarrow{\delta_3} Z_3 \xleftarrow{\delta_4} Z_4 \xleftarrow{\delta_5} Z_5 \xleftarrow{\delta_6} \cdots,$$

where

$$\overline{\mu}(1 \otimes 1) = 1, \quad Z_n = \bigoplus_{r+s=n} Z_{rs} \quad \text{and} \quad \delta_n = \sum_{\substack{r+s=n \\ r+l>0}} \sum_{l=0}^s \delta_{rs}^l,$$

is a projective resolution of the E -bimodule E . Moreover, the family of maps

$$\Gamma_*: Z_* \rightarrow X_*,$$

given by

$$\Gamma(1 \otimes \mathbf{x}_{1s} \otimes \mathbf{v}_{1r} \otimes 1) = \sum_{\sigma \in \mathfrak{S}_r} \text{sg}(\sigma) \otimes \mathbf{x}_{1s} \otimes v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(r)} \otimes 1,$$

defines an morphism of E -bimodule complexes from (Z_*, δ_*) to (X_*, d_*) .

Proof. It is clear that each Z_n is a projective E -bimodule and a direct computation shows that Γ_* is a morphism of complexes. Let

$$G_*^0 \subseteq G_*^1 \subseteq G_*^2 \subseteq G_*^3 \subseteq \dots \quad \text{and} \quad F_*^0 \subseteq F_*^1 \subseteq F_*^2 \subseteq F_*^3 \subseteq \dots$$

be the filtration of Z_* and X_* respectively, defined by

$$G_n^i = \bigoplus_{\substack{r+s=n \\ s \leq i}} Z_{rs} \quad \text{and} \quad F_n^i = \bigoplus_{\substack{r+s=n \\ s \leq i}} X_{rs}.$$

In order to see that Γ_* is a quasi-isomorphism it is sufficient to show that it induces a quasi-isomorphism between the graded complexes associated with the filtrations introduced above. In other words that the maps

$$\Gamma_{*s} : (Z_{*s}, \delta_{*s}^0) \longrightarrow (X_{*s}, d_{*s}^0) \quad (s \geq 0),$$

defined by

$$\Gamma(1 \otimes \mathbf{x}_{1s} \otimes \mathbf{v}_{1r} \otimes 1) = \sum_{\sigma \in \mathfrak{S}_r} \text{sg}(\sigma) \otimes \mathbf{x}_{1s} \otimes v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(r)} \otimes 1,$$

are quasi-isomorphisms, which follows easily from Proposition 2.1. $\square$

5.1. Hochschild cohomology. Let M be an E -bimodule. For $r, s \geq 0$, let

$$\overline{Z}^{rs}(M) = \text{Hom}_k(V^r \otimes \mathfrak{g}^{\wedge s}, M).$$

We define the morphism

$$\overline{\delta}_l^{rs} : \overline{Z}^{r+l-1, s-l}(M) \longrightarrow \overline{Z}^{rs}(M) \quad (\text{with } 0 \leq l \leq \min(2, s) \text{ and } r+l > 0)$$

by:

$$\begin{aligned} \overline{\delta}_0(\varphi)(\mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) &= \sum_{i=1}^r (-1)^i [v_i, \varphi(\mathbf{v}_{1\widehat{i}r} \otimes \mathbf{x}_{1s})], \\ \overline{\delta}_1(\varphi)(\mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) &= \sum_{i=1}^s (-1)^{i+r} [\varphi(\mathbf{v}_{1r} \otimes \mathbf{x}_{1\widehat{i}s}), 1 \# x_i] \\ &\quad + \sum_{\substack{i=1 \\ 1 \leq h \leq r}}^s (-1)^{i+r} \varphi(\mathbf{v}_{1, h-1} \wedge v_h^{\overline{x}_i} \wedge \mathbf{v}_{h+1, r} \otimes \mathbf{x}_{1\widehat{i}s}) \\ &\quad + \sum_{1 \leq i < j \leq s} (-1)^{i+j+r} \varphi(\mathbf{v}_{1r} \otimes [x_i, x_j]_{\mathfrak{g}} \wedge \mathbf{x}_{1\widehat{i}\widehat{j}s}) \end{aligned}$$

and

$$\overline{\delta}_2(\varphi)(\mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) = \sum_{1 \leq i < j \leq s} (-1)^{i+j} \varphi(\widehat{f}_{ij} \wedge \mathbf{v}_{1r} \otimes \mathbf{x}_{1\widehat{i}\widehat{j}s}).$$

Applying the functor $\text{Hom}_{E^e}(-, M)$ to the complex (Z_*, δ_*) , and using Theorem 5.1 and the identifications $\xi^{rs} : \overline{Z}^{rs}(M) \rightarrow \text{Hom}_{E^e}(Z_{rs}, M)$, given by

$$\xi(\varphi)(1 \otimes \mathbf{x}_{1s} \otimes \mathbf{v}_{1r} \otimes 1) = (-1)^{rs} \varphi(\mathbf{v}_{1r} \otimes \mathbf{x}_{1s}),$$

we obtain the complex

$$\overline{Z}^0(M) \xrightarrow{\overline{\delta}^1} \overline{Z}^1(M) \xrightarrow{\overline{\delta}^2} \overline{Z}^2(M) \xrightarrow{\overline{\delta}^3} \overline{Z}^3(M) \xrightarrow{\overline{\delta}^4} \overline{Z}^4(M) \xrightarrow{\overline{\delta}^5} \dots,$$

where

$$\overline{Z}^n(M) = \bigoplus_{r+s=n} \overline{Z}^{rs}(M) \quad \text{and} \quad \overline{\delta}^n = \sum_{\substack{r+s=n \\ r+l>0}} \sum_{l=0}^{\min(s,2)} \overline{\delta}_l^{rs}.$$

Note that if $f(\mathfrak{g} \otimes \mathfrak{g}) \subseteq k$, then $(\overline{Z}^*(M), \overline{\delta}^*)$ is the total complex of the double complex $(\overline{Z}^{**}(M), \overline{\delta}_0^{**}, \overline{\delta}_1^{**})$.

Theorem 5.2. *The Hochschild cohomology $H^*(E, M)$, of E with coefficients in M , is the cohomology of $(\overline{Z}^*(M), \overline{\delta}^*)$.*

The map $\Gamma: (Z_*, \delta_*) \rightarrow (X_*, d_*)$ induces a quasi-isomorphism

$$\overline{\Gamma}^*: (\overline{X}_k^*(M), \overline{d}_*) \longrightarrow (\overline{Z}^*(M), \overline{\delta}^*).$$

Proposition 5.3. *We have*

$$\overline{\Gamma}(\varphi)(\mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) = \sum_{\sigma \in \mathfrak{S}_r} \text{sg}(\sigma) \varphi(v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(r)} \otimes \mathbf{x}_{1s}).$$

Proof. This follows immediately from Theorem 5.1. $\square$

5.2. The cup product. In this subsection we compute the cup product of $\text{HH}^*(E)$ in terms of the complex $(\overline{Z}^*(E), \overline{\delta}^*)$. Given $\phi \in \overline{Z}^{rs}(E)$ and $\phi' \in \overline{Z}^{r's'}(E)$, we define $\phi \star \phi' \in \overline{Z}^{r+r', s+s'}(E)$ by

$$(\phi \star \phi')(\mathbf{v}_{1r''} \otimes \mathbf{x}_{1s''}) = \sum_{\substack{1 \leq i_1 < \cdots < i_r \leq r'' \\ 1 \leq j_1 < \cdots < j_s \leq s''}} \text{sg}(i_1, j_1) \phi(\mathbf{v}_{i_1} \otimes \cdots \otimes \mathbf{x}_{j_s}) \phi'(\mathbf{v}_{h_{1r'}} \otimes \mathbf{x}_{l_{1s'}}),$$

where

- $\text{sg}(i_1, j_1) = (-1)^{r's + \sum_{u=1}^r (i_u - u) + \sum_{u=1}^s (j_u - u)}$,
- $r'' = r + r'$ and $s'' = s + s'$,
- $1 \leq h_1 < \cdots < h_{r'} \leq r''$ denote the set defined by
$$\{i_1, \dots, i_r\} \cup \{h_1, \dots, h_{r'}\} = \{1, \dots, r''\},$$
- $1 \leq l_1 < \cdots < l_{s'} \leq s''$ denote the set defined by
$$\{j_1, \dots, j_s\} \cup \{l_1, \dots, l_{s'}\} = \{1, \dots, s''\},$$
- $\mathbf{v}_{i_1} = v_{i_1} \wedge \cdots \wedge v_{i_r}$ and $\mathbf{v}_{h_{1r'}} = v_{h_1} \wedge \cdots \wedge v_{h_{r'}}$,
- $\mathbf{x}_{j_1} = x_{j_1} \wedge \cdots \wedge x_{j_s}$ and $\mathbf{x}_{l_{1s'}} = x_{l_1} \wedge \cdots \wedge x_{l_{s'}}$.

Theorem 5.4. *The cup product of $\text{HH}^*(E)$ is induced by the operation $\star$ in the complex $(\overline{Z}^*(E), \overline{\delta}^*)$.*

Proof. By Theorem 3.4 it suffices to prove that

$$(2) \quad \overline{\Gamma}(\varphi \bullet \varphi') = \overline{\Gamma}(\varphi) \star \overline{\Gamma}(\varphi')$$

for all $\varphi \in \overline{X}_k^{rs}(E)$ and $\varphi' \in \overline{X}_k^{r's'}(E)$. Let $\phi = \overline{\Gamma}(\varphi)$ and $\phi' = \overline{\Gamma}(\varphi')$. On one hand

$$\begin{aligned} (\phi \star \phi')(\mathbf{v}_{1r''} \otimes \mathbf{x}_{1s''}) &= \sum_{\substack{1 \leq i_1 < \cdots < i_r \leq r'' \\ 1 \leq j_1 < \cdots < j_s \leq s''}} \text{sg}(i_1, j_1) \phi(\mathbf{v}_{i_1} \otimes \cdots \otimes \mathbf{x}_{j_s}) \phi'(\mathbf{v}_{h_{1r'}} \otimes \mathbf{x}_{l_{1s'}}) \\ &= \sum_{\substack{1 \leq i_1 < \cdots < i_r \leq r'' \\ 1 \leq j_1 < \cdots < j_s \leq s'' \\ \tau \in \mathfrak{S}_r, \nu \in \mathfrak{S}_{r'}}} \text{sg}(i_1, j_1) \text{sg}(\tau) \text{sg}(\nu) \varphi(v_{i_{\tau(1r)}} \otimes \cdots \otimes \mathbf{x}_{j_{1s}}) \varphi'(\mathbf{v}_{h_{\nu(1r')}} \otimes \mathbf{x}_{l_{1s'}}), \end{aligned}$$

where

$$\mathbf{v}_{i_{\tau(1r)}} = v_{i_{\tau(1)}} \otimes \cdots \otimes v_{i_{\tau(r)}} \quad \text{and} \quad \mathbf{v}_{h_{\nu(1r')}} = v_{h_{\nu(1)}} \otimes \cdots \otimes v_{h_{\nu(r')}}.$$

On the other hand

$$\begin{aligned} \bar{\Gamma}(\varphi \bullet \varphi')(\mathbf{v}_{1r''} \otimes \mathbf{x}_{1s''}) &= \sum_{\sigma \in \mathfrak{S}_{r''}} \text{sg}(\sigma)(\varphi \bullet \varphi')(v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(r'')} \otimes \mathbf{x}_{1s}) \\ &= \sum_{\substack{1 \leq j_1 < \cdots < j_s \leq s'' \\ \sigma \in \mathfrak{S}_{r''}}} \text{sg}(\sigma) \text{sg}(j_s) \varphi_1(v_{\sigma(1r)} \otimes \mathbf{x}_{j_{1s}}) \varphi_2(v_{\sigma(r+1, r'')} \otimes \mathbf{x}_{l_{1s}}), \end{aligned}$$

where

$$v_{\sigma(1r)} = v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(r)} \quad \text{and} \quad v_{\sigma(r+1, r'')} = v_{\sigma(r+1)} \otimes \cdots \otimes v_{\sigma(r'')}.$$

Now, formula (2) if follows immediately from these facts. $\square$

5.3. Hochschild homology. Let M be an E -bimodule. For $r, s \geq 0$, let

$$\bar{Z}_{rs}(M) = M \otimes V^{\wedge r} \otimes \mathfrak{g}^{\wedge s}.$$

We define the morphisms

$$\bar{\delta}_{rs}^l: \bar{Z}_{rs}(M) \longrightarrow \bar{Z}_{r+l-1, s-l}(M) \quad (0 \leq l \leq s \text{ and } r+l > 0)$$

by:

$$\begin{aligned} \bar{\delta}_{rs}^0(m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) &= \sum_{i=1}^r (-1)^i [m, v_i] \otimes \mathbf{v}_{1\widehat{ir}} \otimes \mathbf{x}_{1s}, \\ \bar{\delta}_{rs}^1(m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) &= \sum_{i=1}^s (-1)^{i+r} [1 \# x_i, m] \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1\widehat{is}} \\ &\quad + \sum_{\substack{i=1 \\ 1 \leq h \leq r}}^s (-1)^{i+r} m \otimes \mathbf{v}_{1, h-1} \wedge v_h^{\bar{x}_i} \wedge \mathbf{v}_{h+1, r} \otimes \mathbf{x}_{1\widehat{is}} \\ &\quad + \sum_{1 \leq i < j \leq s} (-1)^{i+j+r} m \otimes \mathbf{v}_{1r} \otimes [x_i, x_j]_{\mathfrak{g}} \wedge \mathbf{x}_{1\widehat{ij}s} \end{aligned}$$

and

$$\bar{\delta}_{rs}^2(m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) = \sum_{1 \leq i < j \leq s} (-1)^{i+j} m \otimes \widehat{f}_{ij} \wedge \mathbf{v}_{1r} \otimes \mathbf{x}_{1\widehat{ij}s}.$$

By tensoring on the left the complex (Z_*, δ_*) over E^e with M , and using Theorem 5.1 and the identifications $\xi_{rs}: \bar{Z}_{rs}(M) \rightarrow M \otimes_{E^e} Z_{rs}$, given by

$$\xi(m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) = (-1)^{rs} m \otimes_{E^e} (1 \otimes \mathbf{x}_{1s} \otimes \mathbf{v}_{1r} \otimes 1),$$

we obtain the complex

$$\bar{Z}_0(M) \xleftarrow{\bar{\delta}_1} \bar{Z}_1(M) \xleftarrow{\bar{\delta}_2} \bar{Z}_2(M) \xleftarrow{\bar{\delta}_3} \bar{Z}_3(M) \xleftarrow{\bar{\delta}_4} \bar{Z}_4(M) \xleftarrow{\bar{\delta}_5} \cdots,$$

where

$$\bar{Z}_n(M) = \bigoplus_{r+s=n} \bar{Z}_{rs}(M) \quad \text{and} \quad \bar{\delta}_n = \sum_{\substack{r+s=n \\ r+l>0}} \sum_{l=0}^{\min(s,2)} \bar{\delta}_{rs}^l.$$

Note that if $f(\mathfrak{g} \otimes \mathfrak{g}) \subseteq k$, then $(\bar{Z}_*(M), \bar{\delta}_*)$ is the total complex of the double complex $(\bar{Z}_{**}(M), \bar{\delta}_{**}^0, \bar{\delta}_{**}^1)$.

Theorem 5.5. *The Hochschild homology $H_*(E, M)$, of E with coefficients in M , is the homology of $(\bar{Z}_*(M), \bar{\delta}_*)$.*

Proof. It is an immediate consequence of the above discussion. $\square$

The map $\Gamma: (Z_*, \delta_*) \rightarrow (X_*, d_*)$ induces a quasi-isomorphism

$$\bar{\Gamma}_*: (\bar{Z}_*(M), \bar{\delta}_*) \longrightarrow (\bar{X}_*^k(M), \bar{d}_*) .$$

Proposition 5.6. *We have*

$$\bar{\Gamma}(m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) = \sum_{\sigma \in \mathfrak{S}_r} \text{sg}(\sigma) m \otimes v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(r)} \otimes \mathbf{x}_{1s} .$$

Proof. This follows immediately from Theorem 5.1. $\square$

5.4. The cap product. In this subsection we compute the cap product of

$$H_p(E, M) \times \text{HH}^q(E) \rightarrow H_{p-q}(E, M) \quad (q \leq p),$$

in terms of the complexes $(\bar{Z}_*(M), \bar{\delta}_*)$ and $(\bar{Z}^*(E), \bar{\delta}^*)$. Given

$$m \otimes \mathbf{v}_{1s} \otimes \mathbf{x}_{1s} \in \bar{Z}_{rs}(M) \quad \text{and} \quad \phi' \in \bar{Z}^{r's'}(E) \quad \text{with } r \geq r' \text{ and } s \geq s',$$

we define $(m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) \star \phi' \in \bar{Z}_{r-r', s-s'}(M)$ by

$$(m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) \star \phi' = \sum_{\substack{1 \leq i_1 < \cdots < i_{r'} \leq r \\ 1 \leq j_1 < \cdots < j_{s'} \leq s}} \text{sg}(i_{1r'}, j_{1s'}) m \phi'(\mathbf{v}_{i_{1r'}} \otimes \mathbf{x}_{j_{1s'}}) \otimes \mathbf{v}_{h_{1, r-r'}} \otimes \mathbf{x}_{l_{1, s-s'}} .$$

where

- $\text{sg}(i_{1r'}, j_{1s'}) = (-1)^{rs' + r's' + \sum_{u=1}^{r'} (i_u - u) + \sum_{u=1}^{s'} (j_u - u)}$,
- $1 \leq h_1 < \cdots < h_{r-r'} \leq r$ denote the set defined by
$$\{i_1, \dots, i_{r'}\} \cup \{h_1, \dots, h_{r-r'}\} = \{1, \dots, r\},$$
- $1 \leq l_1 < \cdots < l_{s-s'} \leq s$ denote the set defined by
$$\{j_1, \dots, j_{s'}\} \cup \{l_1, \dots, l_{s-s'}\} = \{1, \dots, s\},$$
- $\mathbf{v}_{i_{1r'}} = v_{i_1} \wedge \cdots \wedge v_{i_{r'}}$ and $\mathbf{v}_{h_{1, r-r'}} = v_{h_1} \wedge \cdots \wedge v_{h_{r-r'}}$,
- $\mathbf{x}_{j_{1s'}} = x_{j_1} \wedge \cdots \wedge x_{j_{s'}}$ and $\mathbf{x}_{l_{1, s-s'}} = x_{l_1} \wedge \cdots \wedge x_{l_{s-s'}}$.

Theorem 5.7. *The cap product*

$$H_p(E, M) \times \text{HH}^q(E) \rightarrow H_{p-q}(E, M) \quad (q \leq p),$$

is induced by $\star$, in terms of the complexes $(\bar{Z}_*(M), \bar{\delta}_*)$ and $(\bar{Z}^*(E), \bar{\delta}^*)$.

Proof. By Theorem 4.4 it suffices to prove that

$$(3) \quad \bar{\Gamma}(m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) \bullet \phi' = \bar{\Gamma}((m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) \star \bar{\Gamma}(\phi'))$$

for all $m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s} \in \bar{Z}_{rs}(M)$ and $\phi' \in \bar{X}_k^{r's'}(E)$. Let $\phi' = \bar{\Gamma}(\phi')$. On one hand

$$\bar{\Gamma}(m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) \bullet \phi' = \sum_{\substack{1 \leq j_1 < \cdots < j_{s'} \leq s \\ \sigma \in \mathfrak{S}_r}} \text{sg}(\sigma j_{1s'}) m \phi'(\mathbf{v}_{\sigma(1r')} \otimes \mathbf{x}_{j_{1s'}}) \otimes v_{\sigma(r'+1, r)} \otimes \mathbf{x}_{l_{1, s-s'}},$$

where $\text{sg}(\sigma j_{1s'}) = \text{sg}(\sigma) \text{sg}(j_{1s'})$,

$$v_{\sigma(1r')} = v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(r')} \quad \text{and} \quad v_{\sigma(r'+1, r)} = v_{\sigma(r'+1)} \otimes \cdots \otimes v_{\sigma(r)} .$$

On the other hand

$$(m \otimes \mathbf{v}_{1r} \otimes \mathbf{x}_{1s}) \star \phi' = \sum_{\substack{1 \leq i_1 < \cdots < i_{r'} \leq r \\ 1 \leq j_1 < \cdots < j_{s'} \leq s}} \text{sg}(i_{1r'}, j_{1s'}) m \phi'(\mathbf{v}_{i_{1r'}} \otimes \mathbf{x}_{j_{1s'}}) \otimes \mathbf{v}_{h_{1, r-r'}} \otimes \mathbf{x}_{l_{1, s-s'}}$$

$$= \sum_{\substack{1 \leq i_1 < \dots < i_{r'} \leq r \\ 1 \leq j_1 < \dots < j_{s'} \leq s \\ \tau \in \mathfrak{S}_{r'}}} \text{sg}(\tau) \text{sg}(i_{1r'}, j_{1s'}) m\varphi'(\mathbf{v}_{i_{\tau(1r')}} \otimes \mathbf{x}_{j_{1s'}}) \otimes \mathbf{v}_{h_{1, r' - r}} \otimes \mathbf{x}_{l_{1, s' - s}},$$

where $\mathbf{v}_{i_{\tau(1r')}} = v_{i_{\tau(1)}} \otimes \dots \otimes v_{i_{\tau(r')}} \cdot$ Now, formula (3) if follows immediately from these facts. $\square$

REFERENCES

- [B-C-M] R. J. Blattner, M. Cohen and S. Montgomery, *Crossed products and inner actions of Hopf algebras*, Trans. Amer. Math. Soc. 298 (1986) 671–711.
- [B-G-R] W. Borho, P. Gabriel and R. Rentschler, *Primideale in Einhüllenden auflösbarer Lie-Algebren*, Lecture Notes in Mathematics 357, Springer-Verlag, Berlin Heidelberg New York (1973).
- [Ch] W. Chin, *Prime ideals in differential operator rings and crossed products of infinite groups*, Journal of Algebra 106 (1987) 78–104.
- [D-T] Y. Doi and M. Takeuchi, *Cleft comodule algebras by a bialgebra*, Comm. in Alg. 14 (1986) 801–817.
- [G-G1] J.A. Guccione and J.J. Guccione, *Hochschild (co)homology of differential operators rings*, Journal of Algebra 243 (2001) 596–614.
- [G-G2] J.A. Guccione and J.J. Guccione, *Hochschild (co)homology of a Hopf crossed products*, K-theory 25 (2002) 139–169.
- [Mc] J. C. McConnell, *Representations of solvable Lie algebras and the Gelfand-Kirillov conjecture*, Proc. LMS 29 (1974) 453–484.
- [Mc-R] J. C. McConnell and J. C. Robson, *No commutative Noetherian rings*, Wiley-Interscience New York (1987).
- [M] S. Montgomery, *Crossed products of Hopf algebras and enveloping algebras*, Perspectives in ring theory, Kluwer Academic, Dordrecht (1988) 253–268.
- [S] R. Sridharan, *Filtered algebras and representations of Lie algebras*, Trans. Am. Math. Soc. 100 (1961) 530–550.

DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS Y NATURALES, PABELLÓN 1 - CIUDAD UNIVERSITARIA, (1428) BUENOS AIRES, ARGENTINA.

DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS Y NATURALES, PABELLÓN 1 - CIUDAD UNIVERSITARIA, (1428) BUENOS AIRES, ARGENTINA.
E-mail address: **vander@dm.uba.ar**

DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS Y NATURALES, PABELLÓN 1 - CIUDAD UNIVERSITARIA, (1428) BUENOS AIRES, ARGENTINA.
E-mail address: **jgucci@dm.uba.ar**